

Order: order@swbio.com



Catalog Number: 21066-1, 21066-2 **Amount:** 50µg/50µl, 100µg/100µl

Swiss-Prot No.: P03372

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl,0.02% sodium azide and 50% glycerol. **Storage/Stability:** Store at -20°C/1 year

Immunogen: The antiserum was produced against synthesized non-phosphopeptide derived from

Human Estrogen Receptor-α around the phosphorylation site of serine 106 (S-P- SP-P-L).

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen


Specificity/Sensitivity:Estrogen Receptor-α (Ab-106) antibody detects endogenous levels of total Estrogen Receptor-α protein.

Reactivity: Human, Mouse

Applications:

Predicted MW: 66 kd

WB: 1:500~1:1000 IHC:1:50~1:100




Estradiol - - - +
Peptide - + - -

Western blot analysis of extracts from MCF7 cells using

Estrogen Receptor-α (Ab-106) antibody (#21066) and

Estrogen Receptor-α (phospho-Ser106) antibody (#11071).



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Estrogen Receptor-α (Ab-106) antibody (#21066).

Background:

Nuclear hormone receptor. The steroid hormones and their receptors are involved in the regulation of eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues.

References:

H Yamaguchi, J Zhu, T Yu, et al. (2007) Serum-free mouse embryo cells generate a selfsustaining feedback loop for an astrocyte marker protein and respond to cytokines and bisphenol A in accordance with the subtle difference in their differentiation state. Cell Biology International, 31(6):638-644.