



P53 (Ab-33)  
Antibody

#21088

**Catalog Number:** 21088-1, 21088-2

**Amount:** 50µg/50µl, 100µg/100µl

**Swiss-Prot No. :** P04637

**Form of Antibody:** Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

**Storage/Stability:** Store at -20°C/1 year

**Immunogen:** The antiserum was produced against synthesized non-phosphopeptide derived from human p53 around the phosphorylation site of serine serine33 (V-L-S<sup>P</sup>-P-L).

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen


**Specificity/Sensitivity**p53 (Ab-37) antibody detects endogenous levels of total p53 protein

**Reactivity:** Human,

**Applications:**

Predicted MW: 53 kd

WB: 1:500~1:1000



Western blot analysis of extracts from HT-29 cells untreated or treated with UV (20min), using p53 (Ab-33) antibody (#21088, Lane1 and 2) and p53 (phospho-Ser33) antibody (#11097, Lane 3 and 4).

### Background :

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Implicated in Notch signaling cross-over

### References:

- Lin T, et al. (2005) Nat Cell Biol; 7(2): 165-71.
- Vega FM, et al. (2004) Mol Cell Biol; 24(23): 10366-80.
- Li J, et al. (2004) J Biol Chem; 279(40): 41275-9.
- Wang J, et al. (2004) J Biol Chem; 279(38): 39584-92.