



P53 (Ab-315)
Antibody

#21091

Catalog Number: 21091-1, 21091-2

Amount: 50µg/50µl, 100µg/100µl

Swiss-Prot No. : P04637

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Storage/Stability: Store at -20°C/1 year

Immunogen: The antiserum was produced against synthesized non-phosphopeptide derived from human p53 around the phosphorylation site of serine 315 (S-S-S_P-P-Q).

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen


Specificity/Sensitivity p53 (Ab-315) antibody detects endogenous levels of total p53 protein

Reactivity: Human,

Applications:

Predicted MW: 53 kd

WB: 1:500~1:1000



Western blot analysis using p53 (Ab-315) antibody (#21091).

Lane 1: HT29 cells; Lane 2: MDA-MB-435 cells;

Lane 3: K562 cells; Lane 4: COLO205 cells.

Background :

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Implicated in Notch signaling cross-over

References:

Lu, H. et al. (1997) Mol. Cell. Biol. 17, 5923-5934.

Lohrum, M. et al. (1996) Oncogene 13, 2527-2539.

Ulrich, S. J. et al. (1993) Proc. Natl. Acad. Sci. USA 90, 5954-5958.

Pospíšilová S, et al. (2004) Biochem J; 378(Pt 3): 939-47.

Merrick BA, et al. (2001) Biochemistry; 40(13): 4053-66.