

Src (Phospho-Tyr529) Signalway Antibody

Catalog Number: 11153-1, 11153-2 Amount: 50µg/50µl, 100µg/100µl

Swiss-Prot No.: P12931

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl,0.02% sodium azide and 50% glycerol. Storage/Stability: Store at -20°C/1 year

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from

human Src around the phosphorylation site of tyrosine 529 (P-Q-Y^P-Q-P).


Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.

Specificity/Sensitivity: Src (phospho-Tyr529) antibody detects endogenous levels of Src only when phosphorylated at tyrosine 529.

Reactivity: Human, Mouse, Rat

Applications:


Predicted MW: 60 kd




EGF+serum -- + -

Peptide - + - -

Western blot analysis of extracts from 293 cells using Src (Ab-529) antibody (#21168, Lane1 and 2) and Src (phospho-Tyr529) antibody (#11153, Lane 3 and 4).



P-Peptide - +
Immunohistochemical analysis of paraffin- embedded
human lung carcinoma tissue using Src (phospho-Tyr529) antibody (#11153).



Immunofluorescence staining of methanol-fixed HeLa cells using Src (phospho- Tyr529)antibody (#11153, Red).

Background:

proto-oncogenic cytoplasmic tyrosine kinase of the SRC family. Highly expressed in certain fully differentiated cells such as neurons, platelets and macrophages. Phosphorylation of an activation loop tyrosine activates the enzyme; phosphorylation of a tyrosine in the C-terminus by Csk inhibits the enzyme.

References:

.Pyper J.M., (1985) Mol. Cell. Biol. 5:831-838 Pyper J.M.(1990) Mol. Cell. Biol. 10:2035-2040 Pyper J.M.(1990) Mol. Cell. Biol. 10:2035-2040 Xu W., (1997).Nature 385:595-602 Benes C.H., (2005) Cell 121:271-280