



## BAG1 Antibody

#24175

**Catalog Number:** 24175-1, 24175-2

**Amount:** 50µg/50µl, 100µg/100µl

**Swiss-Prot No. :** Q99933

**Form of Antibody:** Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

**Storage/Stability:** Store at -20°C/1 year


**Immunogen:** The antiserum was produced against synthesized peptide derived from Human BAG1

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

**Specificity/Sensitivity:** BAG1 antibody detects endogenous levels of total BAG1 protein

**Reactivity:** Human, Mouse, Rat

**Applications:** Predicted MW: 39kd      WB:1:500~1:2000      IHC:1:50-200



Western blot analysis of extracts of NCI-H460 cell lines, using BAG1 antibody.

**Background :** Bag1 belongs to the Bcl-2 associated athanogene (BAG) family of multifunctional proteins and was the first of six related proteins isolated from humans . This widely expressed protein interacts with a number of signaling molecules (including Bcl2, HGF receptor and Raf1) as it regulates signaling molecules in pathways involving cell survival, growth and differentiation. The most common role played by Bag1 protein is as an inhibitor of proteins favoring apoptosis . Bag1 also plays a role in Raf1 signaling and binds DNA as a transcription activator . Bag1 protein is a well-characterized inhibitor of its binding partner HSP70 . A conserved carboxy-terminal BAG domain within Bag1 interacts with the ATPase domain of HSP70 to negatively regulate heat shock protein chaperone activity . The multiple isoforms of Bag1 protein generated from a single transcript share a common ubiquitin homology domain and a carboxy-terminal Hsp70 binding region but differ in length and cellular localization. The 50 kDa long (Bag1-L) isoform also contains a nuclear localization signal and is often found in the nucleus where it activates transcription. The 46 kDa intermediate (Bag1-M) isoform is found mainly in the cytoplasm but can also translocate to the nucleus when associated with other proteins. The shorter 29-33 kDa isoforms (Bag1-S, Bag-1) isoforms are found primarily in the cytoplasm . High expression of the anti-apoptotic Bag1 protein correlates with increased survival in patients with particular forms of cancer, leading researchers to study possible therapeutic roles for Bag1 protein .