

CTBP1

Order: order@swbio.com



Catalog Number: 24228-1, 24228-2 **Amount:** 50μg/50μl, 100μg/100μl

Swiss-Prot No.: Q13363

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl,0.02% sodium azide and 50% glycerol. Storage/Stability: Store at -20°C/1 year


Immunogen: The antiserum was produced against synthesized peptide derived from Human CTBP1 Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Specificity/Sensitivity: CTBP1 Antibody detects endogenous levels of total CTBP1

Reactivity: Human, Mouse, Rat

Applications:

Predicted MW: 48kd WB:1:500-2000 IHC:1:50-200



Western blot analysis of extracts of various celllines, using CTBPlantibody.

Background :CtBP1 (C-terminal binding protein 1) was first recognized as a cellular factor that interacts with the C-terminal portion of adenovirus E1A, a protein involved in the transcriptional regulation of key cellular genes. CtBP1 is able to regulate gene activity through its intrinsic dehydrogenase activity and by interacting with Polycomb Group (PcG) proteins during development . Along with its homologue, CtBP2, it acts as a transcriptional corepressor of zinc-finger homeodomain factor deltaEF1 to regulate a wide range of cellular processes through transrepression mechanisms. Through its direct interaction with PRDM16, CtBP1 has been shown to be involved in brown adipose tissue differentiation by mediating the repression of white fat genes and directing differentiation toward the brown fat gene program . CtBP1 also plays a role in lipid metabolic pathways and membrane fission by regulating the fission machinery operating Golgi tubular networks. CtBP1 has recently been shown to repress transcription of BRCA1 via a redox regulated mechanism . Furthermore, it is thought that downregulation of BRCA1 and E-cadherin in invasive ductal breast carcinoma correlates directly with activation of CtBP1.