

Catalog Number: 11578-1, 11578-2

Amount: 50µg/50µl, 100µg/100µl

Swiss-Prot No. : P68431

**Form of Antibody:** Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl,0.02% sodium azide and 50% glycerol.

**Storage/Stability:** Store at -20°C/1 year

**Immunogen:** The antiserum was produced against synthesized phosphopeptide derived from human Histone H3 around the phosphorylation site of Ser28


**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.

**Specificity/Sensitivity:** Histone H3 (phospho-Ser28) antibody detects endogenous levels of Histone H3 only when phosphorylated at Ser28.

Reactivity: Human, Mouse, Rat

## Applications:

Predicted MW: 17 kd WB: 1:500~1:1000



Western blot analysis of extracts from Hela cells, untreated (-) or treated

**Background** : Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling