

Catalog Number: 15577-1, 15577-2

Amount: 50µg/50µl, 100µg/100µl

Swiss-Prot No. : P68431

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl,0.02% sodium azide and 50% glycerol.

Storage/Stability: Store at -20°C/1 year

Immunogen: The antiserum was produced against synthesized peptide derived from


human Histone H3 around the acetylation site of Lysine 14

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Specificity/Sensitivity: HistoneH3(Acetyl-Lys14) antibody detects endogenous levels of Histone H3 only when acetylated at Lys14.

Reactivity: Human, Mouse, Rat

Applications:



Western blot analysis of extracts from Hela cells, untreated (-) or treated

Background : Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling